

## RÉFÉRENCES BIBLIOGRAPHIQUES DES ANNEXES 9-1, 9-2 ET 9-3

- Collins, A., Walling, D., Leeks, G. (1997) Use of the geochemical record preserved in floodplain deposits to reconstruct recent changes in river basin sediment sources. *Geomorphology* 19, 151-167.
- Hancock, G.J., Wilkinson, S.N., Hawdon, A.A., Keen, R.J. (2014) Use of fallout tracers  $^7\text{Be}$ ,  $^{210}\text{Pb}$  and  $^{137}\text{Cs}$  to distinguish the form of sub-surface soil erosion delivering sediment to rivers in large catchments. *Hydrol. Process.* 28, 3855-3874.
- Lacey, J.P., Evrard, O., Smith, H.G., Blake, W.H., Olley, J.M., Minella, J.P.G., Owens, P.N. (2017) The challenges and opportunities of addressing particle size effects in sediment source fingerprinting: A review. *Earth-Sci. Rev.* 169, 85-103.
- Le Gall, M., (2016) Traçage des sources de sédiments à l'amont des hydrosystèmes agricoles: apport de la géochimie élémentaire, du rapport  $^{87}\text{Sr}/^{86}\text{Sr}$  et des radionucléides. Paris Saclay.
- Legout, C., Poulenard, J., Nemery, J., Navratil, O., Grangeon, T., Evrard, O., Esteves, M. (2013) Quantifying suspended sediment sources during runoff events in headwater catchments using spectrophotometry. *Journal of Soils and Sediments* 13, 1478-1492.
- Ploquin, F., Fritsch, E., Juillot, F., Dublet, G., Estève, I., Fandeur, D., Fiallin, M., Guinier, J., (2017) Gisements de nickel latéritique, volume IV: les oxydes de manganèse associés aux latérites de transition. CNRT "Nickel et son environnement".
- Schwertmann, U., Latham, M. (1986) Properties of iron oxides in some New Caledonian oxisols. *Geoderma* 39, 105-123.
- Sevin, B., (2014) Cartographie du régolithe sur formation ultrabasique de Nouvelle-Calédonie: Localisation dans l'espace et le temps des gisements nickelifères. Nouvelle Calédonie.
- Wilkinson, S.N., Hancock, G.J., Bartley, R., Hawdon, A.A., Keen, R.J. (2013) Using sediment tracing to assess processes and spatial patterns of erosion in grazed rangelands, Burdekin River basin, Australia. *Agric. Ecosyst. Environ.* 180, 90-102.
- Croudace, I. W., & Rothwell, R. G. (Eds.). (2015). *Micro-XRF Studies of Sediment Cores Applications of a non-destructive tool for the environmental sciences (Vol. 17)*. Springer.
- Foucher, A., Salvador-Blanes, S., Evrard, O., Simonneau, A., Chapron, E., Courp, T., ... & Desmet, M. (2014). Increase in soil erosion after agricultural intensification: evidence from a lowland basin in France. *Anthropocene*, 7, 30-41
- Garcin M., Gastaldi Y., Lesimple S. (2017) Quantification et évolution temporelle des apports miniers dans les rivières calédoniennes. BRGM/RP-66840-FR, 44 p., 23 fig., 5. Bureau des Recherches Géologiques et Minières.
- Le Gall, M., Evrard, O., Foucher, A., Lacey, J. P., Salvador-Blanes, S., Thil, F., Dapoigny, A., Lefèvre, I., Cerdan, O., Ayrault, S. (2016 b). Quantifying sediment sources in a lowland agricultural catchment pond using  $^{137}\text{Cs}$  activities and radiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$ . *Science Total of the Environment* 566, 968-980.

Scheinost, A. C., Chavernas, A., Barrón, V., & Torrent, J. (1998). Use and limitations of second-derivative diffuse reflectance spectroscopy in the visible to near-infrared range to identify and quantify Fe oxide minerals in soils. *Clays and Clay Minerals*, 46(5), 528-536.

Walling D, Peart M, Oldfield F, Thompson R (1979) Suspended sediment sources identified by magnetic measurements. *Nature* 281:110.

Rapport de stage de Lucie Gosset (2016). Rivières engravées à Thio, Nouvelle-Calédonie : vécus et attentes de la population